

A GEOMETRIC PROOF OF THE CLASSIFICATION OF COMPLEX VECTOR CROSS PRODUCT

CHENGJIE YU

ABSTRACT. In this article, we give a geometric proof of the classification of complex vector cross product in Lee-Leung [1].

The following definition is a point-wise version of the definition of complex vector cross product on complex Hermitian manifolds in Lee-Leung [1].

Definition 0.1. Let V be complex vector space of dimension n with a Hermitian metric and r be an integer between 1 and $n - 1$. A $(r + 1)$ -form $\phi \in \wedge^{r+1}V^*$ is called a complex r -fold vector cross product on V if

$$(0.1) \quad \|\iota_{e_1 \wedge e_2 \wedge \cdots \wedge e_r} \phi\|^2 = 1$$

for any r unit vectors $e_1, e_2, \dots, e_r$ that are orthogonal to each other.

In [1], Lee-Leung proved the following result.

Theorem 0.1. *On a complex vector space V of dimension n , there is a r -fold complex vector cross product if and only if*

- (1) $r = 1$ and n is an even number, or
- (2) $r = n - 1$ and n is an arbitrary natural number.

In this article, we give a geometric proof of this theorem by using Plücker embedding and dimension counting of projective sub-varieties.

Proof. The "if" part is clear (Ref. Lee-Leung [1]). We only prove the "only if" part.

Let $\phi \in \wedge^{r+1}V^*$ be a r -fold complex vector product. Then, it induces a complex linear map $f : \wedge^r V \rightarrow V^*$ by sending $u_1 \wedge u_2 \wedge \cdots \wedge u_r$ to $\iota_{u_1 \wedge u_2 \wedge \cdots \wedge u_r} \phi$.

Date: Oct. 2008.

By the identity (0.1), we know that for any $u_1, u_2, \dots, u_r$ that are linearly independent,

$$u_1 \wedge u_2 \wedge \dots \wedge u_r \notin \ker f.$$

By projectification and Plücker embedding, all the r -vectors that can be separated as the form $u_1 \wedge u_2 \wedge \dots \wedge u_r$ forms the Grassmannian manifold $Gr(r, V) \subset \mathbb{P}(\wedge^r V)$. The discussion above implies that that

$$Gr(r, V) \cap \mathbb{P}(\ker f) = \emptyset.$$

Hence (Ref. Shafarevich [2])

$$\dim Gr(r, V) + \dim \mathbb{P}(\ker f) < \dim \mathbb{P}(\wedge^r V).$$

That is,

$$(0.2) \quad \dim Gr(r, V) + \dim \ker f < \dim(\wedge^r V).$$

Note that, by dimension theorem in linear algebra, we have

$$(0.3) \quad \dim \ker f \geq \dim \wedge^r V - n.$$

Combining equation (0.2) and (0.3), we have

$$r(n - r) = \dim Gr(r, V) < n.$$

Therefore $r = 1$ or $r = n - 1$.

When $r = 1$, $f : V \rightarrow V^*$ is an isometry satisfying that

$$\langle f(u), u \rangle = \phi(u, u) = 0.$$

Let $e_1, e_2, \dots, e_n$ be an orthonormal basis of V and $\omega_1, \omega_2, \dots, \omega_n$ be its dual basis. Then, under this two basis, the matrix A of f is skew symmetric and nonsingular. Hence, n is an even number.

□

REFERENCES

- [1] Lee Jae-Hyouk, Leung Naichung Conan. *Instantons and branes in manifolds with vector cross product*, arXiv:math/0402044v1.
- [2] Shafarevich, Igor R. *Basic algebraic geometry. 1. Varieties in projective space*. Second edition. Translated from the 1988 Russian edition and with notes by Miles Reid. Springer-Verlag, Berlin, 1994. xx+303 pp.

DEPARTMENT OF MATHEMATICS, SHANTOU UNIVERSITY, SHANTOU, GUANGDONG, P.R. CHINA

E-mail address: `cjyu@stu.edu.cn`